





6/11/2014


Chestnut Blights

B3 Summer Science Camp at Olympic High School


Dr. Jennifer Weller

Chestnut Blight - #1

- Cryphonectira parasitica is a fungus (Ascomycete) that affects mostly chestnuts, but also some oaks.
 - The fungus forms orange-yellow fruiting bodies that are easy to see.
 - They form a canker on the tree stems (both trunk and branches).
 - They cannot enter the roots.








Drawings: Ron Peterson and Carole Kroger.





Cryphonectria parasitica up close





Chestnut Blight - #2

- Phytophtora (fie-toff-thora) cinnamomi: The Plant Destroyer
 - ~60 species are known
 - A soil fungus that destroys root tissue and stems roots cannot absorb water



Hyphae, the vegetative state of Phytophthora cinnamomi



Sporangium with zoospores, the main reproductive propagule, of *Phytophthora cinnamomi*

Phytophthora blighted trees




Phytophthora up close

Mycelia spores (a,b)

2-celled version (c,d)

 5d colony on 2 types of media (e,f)




Dr. Jennifer Weller

Plants and their pathogens


- In mammals the immune system can adapt in real time to invasion by pathogens
- Plants lack an adaptive immune system their resistance occurs in 3 ways
 - Specific genetic material, the gene-for-gene method
 - A general chemical arsenal (taxol, salicylic acid, etc)
 - The Hypersensitive Response (HR) which is programmed cell death.


Plant-Pathogen Warfare

- Pathogen-Associated Molecular Patterns (PAMPs) lead to PAMP-Triggered Immunity (PTI)
 - PAMPs are proteins in the pathogen that the plant recognizes with specialized receptors (plant-recognition receptors, PPRs)
 - The interaction starts a signalling cascade and the cell responds by thickening the cell wall, producing toxic compounds, etc.
- Pathogens fight back: a type III secretion system in the pathogen will inject proteins into the host that block the PTI.
 - Virulence proteins are called effectors
- Plants fight back: Effector-triggered Immunity in the plant blocks the effector proteins
 - When the Effector-triggered immune protein binds to the effector the result is programmed cell death, the hypersensitive response.




The Signalling Cascade





Plant-Pathogen Warfare - The Hypersensitive Response in pictures



The Gene-for-Gene Model of Plant Immunity


Resistance gene Effector gene	R	r
A	Resistance	Disease
а	Disease	Disease

Bacterium

Effector protein







Plant


Resistance protein





Gene-for-Gene testing

 What type of experiment might you design to test whether a plant has a resistance gene for a particular pathogen?



Gene-for-gene resistance conferred by the A. thaliana RPS4 gene. Leaves were inoculated with Pseudomonas syringae. Disease (chlorosis) occurs when the host resistance gene (left column) or the pathogen avirulence gene (top row) is absent. Resistance only ensues when both genes are present.

Creating resistant plants

- One pathogen (Cryphonectria parasitica) killed 4 billion tress over 50 years, in an area the size of Montana.
 - 1 tree in 5 in the forests died: oak, hickory and birch surviving of the hard woods
- This implies no natural resistance in the trees. What strategies does that leave us?
 - Breeding in natural resistance from related species
 - This mixes genomes and some traits unique to the American Chestnut may be lost
 - Use biotechnological approaches to insert just the resistance genes
 - Infect the fungus with a virus that kills is (hypovirulence)

TACF's Backcross Breeding Program

Each generation select for:

- ***** Blight resistance
- * American characteristics

Final Product: Goal is true-breeding American type with a high level of blight resistance

