

# The spherical Pythagorean theorem

**Proposition 1** *On a sphere of radius  $R$ , any right triangle  $\triangle ABC$  with  $\angle C$  being the right angle satisfies  $\cos(c/R) = \cos(a/R) \cos(b/R)$ .*

**Proof:** We complement the proof presented in [1, page 206]. Let  $O$  be the center of the sphere, we may assume its coordinates are  $(0, 0, 0)$ . We may rotate the sphere so that  $A$  has coordinates  $\overrightarrow{OA} = (R, 0, 0)$  and  $C$  lies in the  $xy$ -plane. Rotating around the  $z$  axis by  $\beta := \angle AOC$  takes  $A$  into  $C$ . The edge  $OA$  moves in the  $xy$ -plane, by  $\beta$ , thus the coordinates of  $C$  are  $\overrightarrow{OC} = (R \cos(\beta), R \sin(\beta), 0)$ . Since we have a right angle at  $C$ , the plane of  $\triangle OBC$  is perpendicular to the plane of  $\triangle OAC$  and it contains the  $z$  axis. An orthonormal basis of the plane of  $\triangle OBC$  is given by  $1/R \cdot \overrightarrow{OC} = (\cos(\beta), \sin(\beta), 0)$  and the vector  $\overrightarrow{OZ} := (0, 0, 1)$ . A rotation around  $O$  in this plane by  $\alpha := \angle BOC$  takes  $C$  into  $B$ :

$$\overrightarrow{OB} = \cos(\alpha) \cdot \overrightarrow{OC} + \sin(\alpha) \cdot R \cdot \overrightarrow{OZ} = (R \cos(\beta) \cos(\alpha), R \sin(\beta) \cos(\alpha), R \sin(\alpha)).$$

Introducing  $\gamma := \angle AOB$ , we have

$$\cos(\gamma) = \frac{\overrightarrow{OA} \cdot \overrightarrow{OB}}{R^2} = \frac{R^2 \cos(\alpha) \cos(\beta)}{R^2}.$$

The statement now follows from  $\alpha = a/R$ ,  $\beta = b/R$  and  $\gamma = c/R$ . ◇

To prove the rest of the formulas of spherical trigonometry, we need to show the following.

**Proposition 2** *Any spherical right triangle  $\triangle ABC$  with  $\angle C$  being the right angle satisfies*

$$\sin(A) = \frac{\sin\left(\frac{a}{R}\right)}{\sin\left(\frac{c}{R}\right)} \quad \text{and} \tag{1}$$

$$\cos(A) = \frac{\tan\left(\frac{b}{R}\right)}{\tan\left(\frac{c}{R}\right)}. \tag{2}$$

**Proof:** We complement the proof presented in [1, page 208]. After replacing  $a/R$ ,  $b/R$  and  $c/R$  with  $a$ ,  $b$ , and  $c$  we may assume  $R = 1$ . This time we rotate the triangle in such a way that  $\overrightarrow{OC} = (0, 0, 1)$ ,  $A$  is in the  $xz$  plane and  $B$  is in the  $yz$ -plane. A rotation around  $O$  in the  $xz$  plane by  $b = \angle AOC$  takes  $C$  into  $A$ , thus we have

$$\overrightarrow{OA} = (\sin(b), 0, \cos(b)).$$

Similarly, a rotation around  $O$  in the  $yz$ -plane by  $a = \angle BOC$  takes  $C$  into  $B$ , thus we have

$$\overrightarrow{OB} = (0, \sin(a), \cos(a)).$$

The angle  $A$  is between  $\overrightarrow{OA} \times \overrightarrow{OB}$  and  $\overrightarrow{OA} \times \overrightarrow{OC}$ . Here

$$\overrightarrow{OA} \times \overrightarrow{OB} = (-\cos(b) \sin(a), -\sin(b) \cos(a), \sin(a) \sin(b)) \quad \text{and} \quad \overrightarrow{OA} \times \overrightarrow{OC} = (0, -\sin(b), 0).$$

The length of  $\vec{OA} \times \vec{OB}$  is  $|\vec{OA}| \cdot |\vec{OB}| \cdot \sin(c) = \sin(c)$ , the length of  $\vec{OA} \times \vec{OC}$  is  $\sin(b)$ .

To prove (1) we use the fact that

$$|(\vec{OA} \times \vec{OB}) \times (\vec{OA} \times \vec{OC})| = |\vec{OA} \times \vec{OB}| \cdot |\vec{OA} \times \vec{OC}| \cdot \sin(A). \quad (3)$$

Since

$$(-\cos(b)\sin(a), -\sin(b)\cos(a), \sin(a)\sin(b)) \times (0, -\sin(b), 0) = (\sin(a)\sin^2(b), 0, \sin(b)\cos(b)\sin(a))$$

the left hand side of (3) is

$$\sqrt{\sin^2(a)\sin^4(b) + \sin^2(b)\cos^2(b)\sin^2(a)} = \sin(b)\sin(a)\sqrt{\sin^2(b) + \cos^2(b)} = \sin(b)\sin(a).$$

The right hand side of (3) is  $\sin(b)\sin(c)\sin(A)$ . Thus we have

$$\sin(b)\sin(a) = \sin(b)\sin(c)\sin(A),$$

yielding (1).

To prove (2) we use the fact that

$$(\vec{OA} \times \vec{OB}) \cdot (\vec{OA} \times \vec{OC}) = |\vec{OA} \times \vec{OB}| \cdot |\vec{OA} \times \vec{OC}| \cdot \cos(A). \quad (4)$$

The left hand side is  $\sin^2(b)\cos(a)$ , the right hand side is  $\sin(b)\sin(c)\cos(A)$ . Thus we obtain

$$\sin^2(b)\cos(a) = \sin(b)\sin(c)\cos(A),$$

yielding

$$\cos(A) = \frac{\sin(b)\cos(a)}{\sin(c)} = \frac{\tan(b)}{\tan(c)} \cdot \frac{\cos(a)\cos(b)}{\cos(c)}.$$

Equation (2) now follows from Proposition 1. ◇

## References

- [1] D. Royster, "Non-Euclidean Geometry and a Little on How We Got There," Lecture notes, May 7, 2012.