

## Inclusion-exclusion formulas

Let  $A_1, A_2, \dots$ , and  $A_n$  be subsets of the same finite set  $X$ . The *inclusion-exclusion formulas* allow us to count the elements in the union  $\bigcup_{i=1}^n A_i$  and in its complement.

**Theorem 1** *We have*

$$\left| X \setminus \bigcup_{i=1}^n A_i \right| = |X| - \sum_{i=1}^n |A_i| + \sum_{1 \leq i < j \leq n} |A_i \cap A_j| - \cdots + (-1)^n |A_1 \cap A_2 \cap \cdots \cap A_n|, \quad \text{that is,}$$

$$\left| X \setminus \bigcup_{i=1}^n A_i \right| = \sum_{r=0}^n (-1)^r \sum_{1 \leq i_1 < i_2 < \cdots < i_r \leq n} |A_{i_1} \cap A_{i_2} \cap \cdots \cap A_{i_r}|.$$

**Proof:** Let  $x$  be any element of  $X$ . Assume  $x$  belongs to exactly  $j$  sets from  $A_1, \dots, A_n$ . (Here  $j$  is any integer between 0 and  $n$ .) Without loss of generality we may assume that  $x$  belongs to  $A_1, A_2, \dots, A_j$ , but does not belong to  $A_{j+1}, A_{j+2}, \dots, A_n$ . This element is counted zero times on the left hand side if  $j > 0$ , and it is counted once if  $j = 0$ . It suffices to show that  $x$  is counted the same number of times on the right hand side. Clearly  $x$  belongs to the intersection  $A_{i_1} \cap A_{i_2} \cap \cdots \cap A_{i_r}$  if and only if the index set  $\{i_1, \dots, i_r\}$  is a subset of  $\{1, 2, \dots, j\}$ , and there are  $\binom{j}{r}$  such subsets. Thus  $x$  is counted with multiplicity  $\sum_{r=0}^j (-1)^r \binom{j}{r}$  on the right hand side. By the binomial theorem, we have

$$\sum_{r=0}^j (-1)^r \binom{j}{r} = (1 - 1)^j = \delta_{0,j}$$

where  $\delta_{0,j}$  is the Kronecker delta. Therefore  $x$  is counted the same number of times on both sides.  $\diamond$

Subtracting both sides of the equation stated in the Theorem above from  $|X|$ , we obtain an important variant of the inclusion-exclusion formula:

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{i=1}^n |A_i| - \sum_{1 \leq i < j \leq n} |A_i \cap A_j| + \cdots + (-1)^{n-1} |A_1 \cap A_2 \cap \cdots \cap A_n|, \quad \text{that is,}$$

$$\left| \bigcup_{i=1}^n A_i \right| = \sum_{r=1}^n (-1)^{r-1} \sum_{1 \leq i_1 < i_2 < \cdots < i_r \leq n} |A_{i_1} \cap A_{i_2} \cap \cdots \cap A_{i_r}|.$$